

BREAKING NEWS: FUKUSHIMA: Formation of a Bacterium Type which Absorbs the Radioactive Substance

They found plutonium and strontium in every sample

By [Global Research](#)

Global Research, June 05, 2012

Dissensus Japan (translated from Japanese)
5 June 2012

Region: [Asia](#)

Theme: [Environment](#)

Minamisōma (南相馬市, *Minamisōma-shi*?) is a city located in the Fukushima area, within the 25 km evacuation zone.

June 3, 2012 [blog post](#) by Minamisoma City Councilor Koichi Oyama translated by [Dissensus Japan](#):

[...]The other day when a member of the Diet visited Minamisoma City, he said that he had a meeting with an official beforehand but he got no information about it.

What happened to the Ministry of the Environment?

I exchanged business card with three officials when they came for an assembly on April 26. I sent emails but they never answered.

This is my mail:

To Mr.Taku Ômura (Chief of Fukushima Office at Environmental Restoration of The Ministry of the Environmental), Mr. Yukiharu Kouso (Chief of Branch Office at Hamadori of Environmental Resotation of The Ministry of the Environmental), Mr. Aha (Chief of Branch Office at Hamadori of Environmental Resotation of The Ministry of the Environmental)

Well done for the committee meeting of Mimanisoma city assembly yesterday.

I'm Koichi Oyama, an official of Minamisoma City.

I send you additional informations about the black substance we talked about yesterday.

There are a bacterium type which likes to absorb the radioactive substance in exuberance everywhere on the paved surfaces in the residential area and at residential houses and they drifts on the side walks, on parkings and on the road.

Cesium for example is highly radioactive and yesterday i saw the

result of analysis of Japan Chemical Analysis Center which contain fluid, they found plutonium and strontium in every samples.

[...]

The informations of city center detected by Minamisoma City
Total value of cesium 134 and 137
Bq/kg-dry

- 1,320,000 Haramachi District, Ushigoe algae types
- 1,960,000 Kashimaku District, Jisabara algae types
- 5,570,000 Odaka District, Kanaya algae types
- 16,200 Haramachi District, Kunimi soil
- 793,000 Odata District, Kamimachi algae types
- 430,000 Odata District, Motomachi algae types
- 583,000 Odata District, Motomachi algae types
- 2,970,000 Odata District, Kanaya algae types / soil
- 522,000 Odata District, Kanaya cattle dung

[...]

The original source of this article is Dissensus Japan (translated from Japanese)
Copyright © [Global Research](#), Dissensus Japan (translated from Japanese), 2012

[Comment on Global Research Articles on our Facebook page](#)

[Become a Member of Global Research](#)

Articles by: [Global Research](#)

Disclaimer: The contents of this article are of sole responsibility of the author(s). The Centre for Research on Globalization will not be responsible for any inaccurate or incorrect statement in this article. The Centre of Research on Globalization grants permission to cross-post Global Research articles on community internet sites as long the source and copyright are acknowledged together with a hyperlink to the original Global Research article. For publication of Global Research articles in print or other forms including commercial internet sites, contact: publications@globalresearch.ca

www.globalresearch.ca contains copyrighted material the use of which has not always been specifically authorized by the copyright owner. We are making such material available to our readers under the provisions of "fair use" in an effort to advance a better understanding of political, economic and social issues. The material on this site is distributed without profit to those who have expressed a prior interest in receiving it for research and educational purposes. If you wish to use copyrighted material for purposes other than "fair use" you must request permission from the copyright owner.

For media inquiries: publications@globalresearch.ca