

# Does this CDC Report Actually Show that Mask Wearers Are 17x More Likely to Get COVID-19?

By [John C. A. Manley](#)

Global Research, October 21, 2020

Region: [USA](#)

Theme: [Intelligence](#), [Science and Medicine](#)

*I'm afraid I'm going to be the fact-checking party, again. I must call foul on those claiming that a recent [CDC study](#) shows mask wearers are more likely to get COVID-19 than non-mask wearers.*

*I hate face masks, but I love truth. And, while there is other evidence that does support the proposition that face masks would [increase infection rates](#), this new study, in my estimation, is not one of them.*

Here's exactly what the study is saying:

## **Reported use of cloth face covering or mask 14 days before illness onset (missing = 2)**

Never	6 (3.9)
Rarely	6 (3.9)
Sometimes	11 (7.2)
Often	22 (14.4)
Always	108 (70.6)

The numbers in brackets are percentages of the total of 154 patients diagnosed with SARS-COV-2 in the sample group. What's immediately apparent is an inverse relation between face mask use and the number of patients infected. At first glance, this makes it look like wearing a face mask will increase your chances of being infected with SARS-COV-2 by 17-fold; with 70% of patients always wearing their mask in public, compared to the 4% who never don the muzzle.

That's a 1700% increase in viral infection from wearing face masks. That's way too high! We'd have masked corpses all over the place. While wearing a damp cloth over your nose may certainly increase infection rates, is it not far-fetched to suggest such a dramatic increase? For example, the best study I've seen only showed a [50% increase](#) in bacterial infections.

There are other ways the correlation demonstrated in this study could be explained.



Please note: This report has been corrected. An erratum has been published.

Morbidity and Mortality Weekly Report

### Community and Close Contact Exposures Associated with COVID-19 Among Symptomatic Adults $\geq 18$ Years in 11 Outpatient Health Care Facilities — United States, July 2020

Kira A. Fisher, PhD<sup>1</sup>; Mark W. Tenforde, MD, PhD<sup>2,3</sup>; Loren K. Feldstein, PhD<sup>4</sup>; Christopher J. Lindsell, PhD<sup>5,6</sup>; Nathan I. Shapiro, MD<sup>5,7</sup>; D. Clark Fries, MD<sup>5,8</sup>; Kevin W. Gibbs, MD<sup>5,9</sup>; Heidi L. Enderson, MD<sup>10</sup>; Matthew E. Prekker, MD<sup>11</sup>; Jay S. Strengers, MD<sup>12</sup>; Matthew C. Eklund, MD<sup>13</sup>; David T. Henning, MD<sup>14</sup>; Jonathan G. Wilson, MD<sup>15</sup>; Samuel M. Broyles, MEd<sup>16</sup>; Alan D. Peltan, MD<sup>17</sup>; Todd W. Rice, MD<sup>18</sup>; David N. Hahn, MD, PhD<sup>19</sup>; Adri A. Ginde, MD<sup>20</sup>; H. Kristy Talbot, MD<sup>21</sup>; Jonathan D. Casey, MD<sup>22</sup>; Carlos G. Grais, MD<sup>23</sup>; Brendan Flannery, PhD<sup>1</sup>; Mariah M. Patel, MD<sup>1</sup>; Wesley H. Self, MD<sup>24</sup>; IVY Network Investigators; CDC COVID-19 Response Team

Community and close contact exposures continue to drive the coronavirus disease 2019 (COVID-19) pandemic. CDC and other public health authorities recommend community mitigation strategies to reduce transmission of SARS-CoV-2, the virus that causes COVID-19 (1,2). Characterization of community exposures can be difficult to assess when widespread transmission is occurring, especially from asymptomatic persons within inherently interconnected communities. Potential exposures, such as close contact with a person with confirmed COVID-19, have primarily been assessed among COVID-19 cases, without a non-COVID-19 comparison group (3,4). To assess community and close contact exposures associated with COVID-19, exposures reported by case-patients (154) were compared with exposures reported by control-participants (160). Case-patients were symptomatic adults (persons aged  $\geq 18$  years) with SARS-CoV-2 infection confirmed by reverse transcription-polymerase chain reaction (RT-PCR) testing. Control-participants were symptomatic outpatient adults from the same health care facilities who had negative SARS-CoV-2 test results. Close contact with a person with known COVID-19 was more commonly reported among case-patients (42%) than among control-participants (14%). Case-patients were more likely to

This investigation included adults aged  $\geq 18$  years who received a first test for SARS-CoV-2 infection at an outpatient testing or health care center at one of 11 Influenza Vaccine Effectiveness in the Critically Ill (IVY) Network sites\* during July 1–29, 2020 (5). A COVID-19 case was confirmed by RT-PCR testing for SARS-CoV-2 RNA from respiratory specimens. Assays varied among facilities. Each site generated lists of adults tested within the study period by laboratory result; adults with laboratory-confirmed COVID-19 were selected by random sampling as case-patients. For each case-patient, two adults with negative SARS-CoV-2 RT-PCR test results were randomly selected as control-participants and matched by age, sex, and study location. After randomization and matching, 615 potential case-patients and 1,212 control-participants were identified and contacted 14–23 days after the date they received SARS-CoV-2 testing. Screening questions were asked to identify eligible adults. Eligible adults for the study were symptomatic at the time of their first SARS-CoV-2 test. CDC personnel administered structured interviews in English or five other languages<sup>†</sup> by telephone and entered data into REDCap software (6). Among 802 adults contacted and who agreed to participate (295 case-patients and 507

For example, the reason why the majority of the infected patients were frequent face mask users may simply be because the majority of the general population wears masks. The same study found the vast majority of their COVID patient samples were white, non-hispanics. Does that mean white people are more prone to COVID-19? No, it probably just reflects that more white people live in the United States.

“People who are more inclined to wear face masks also may be more inclined to seek out PCR testing and receive positive results,” pointed out [Dr. Ron Brown PhD](#) in an email. “In other words, a limitation of this study, not mentioned by the authors, is selection bias among the sample of ‘314 symptomatic patients who actively sought testing during July 1-29, 2020 at 11 healthcare facilities.’

“Also, a person who is NOT inclined to wear a face mask may be more likely to be a ‘person who did not respond, or refused to participate’ in the study. They are more likely to crawl up in bed and sleep it off.”

I can vouch that if I have a sore throat the last place I’m going is a healthcare facility for a COVID test.

Furthermore, how much can we trust the patients? Imagine your average person in a hospital with a COVID-19 diagnosis: Their doctor asks them if they’ve been wearing their face mask like a good boy or girl. Many who weren’t might lie and say they were.

Amusingly, the study says the researchers used the PCR test to determine whether this sample group had SARS-CoV-2 or not. With all we know of the [PCR test’s shortcomings](#), we might as well say they had the Easter Bunny flip a coin. Why do we fault the PCR test when exposing the corona scandal, but then rely on its junk data to argue that masks are bunk?

In an email, [Prof. Denis Rancourt, PhD](#) offered another perspective on this wonky CDC report:

“The study shows the same proportions [of mask wearing] for the comparison group that did not develop an illness. Therefore, there is no evidence presented in the article that masks increase likelihood of being infected by the presumed virus.”

Thus, we need to be careful about making hasty judgements simply to confirm our position on the [ineffectiveness of masking](#). As Prof. Rancourt’s [arch-enemy, David Kyle Johnson PhD](#), reminds us in a [much-flawed pro-masking article](#): “...once you realize that a few pieces of proposed evidence for something is pseudoscience, you are justified in concluding that all such evidence will be similarly flawed.”

In the end, this new study is just [another example](#) of pointless CDC data; probably intended to trigger false claims and confusion. Or maybe it’s just to make it look like the researchers are researching.

What we really need are large randomly selected samples of people who wear face masks religiously like a crucifix and those who disdain the mask like a dirty diaper. Let’s see what percentage of each group gets ill with flu-like symptoms.

Oh, wait. Of course, we already have [those randomized controlled trial studies](#). And they all say the same thing: Masks make no difference in whether people get sick or not. What more proof do we need that mandatory masking is a baseless violation of personal rights and freedoms?

\*

Note to readers: please click the share buttons above or below. Forward this article to your email lists. Crosspost on your blog site, internet forums. etc.

*John C. A. Manley has spent over a decade ghostwriting for medical doctors, as well as naturopaths, chiropractors and Ayurvedic physicians. He publishes the [COVID-19\(84\) Red Pill Briefs](#) – an email-based newsletter dedicated to preventing the governments of the world from using an exaggerated pandemic as an excuse to violate our freedom, health, privacy, livelihood and humanity. He is also writing a novel, [Brave New Normal: A Dystopian Love Story](#). Visit his website at: [MuchAdoAboutCorona.ca](#) He is a frequent contributor to Global Research.*

The original source of this article is Global Research  
Copyright © [John C. A. Manley](#), Global Research, 2020

---

[Comment on Global Research Articles on our Facebook page](#)

[Become a Member of Global Research](#)

Articles by: [John C. A. Manley](#)

**Disclaimer:** The contents of this article are of sole responsibility of the author(s). The Centre for Research on Globalization will

not be responsible for any inaccurate or incorrect statement in this article. The Centre of Research on Globalization grants permission to cross-post Global Research articles on community internet sites as long the source and copyright are acknowledged together with a hyperlink to the original Global Research article. For publication of Global Research articles in print or other forms including commercial internet sites, contact: [publications@globalresearch.ca](mailto:publications@globalresearch.ca)

[www.globalresearch.ca](http://www.globalresearch.ca) contains copyrighted material the use of which has not always been specifically authorized by the copyright owner. We are making such material available to our readers under the provisions of "fair use" in an effort to advance a better understanding of political, economic and social issues. The material on this site is distributed without profit to those who have expressed a prior interest in receiving it for research and educational purposes. If you wish to use copyrighted material for purposes other than "fair use" you must request permission from the copyright owner.

For media inquiries: [publications@globalresearch.ca](mailto:publications@globalresearch.ca)