

The Advent of Extreme Weather Events and Climate Tipping Points

By [Dr. Andrew Glikson](#)

Global Research, March 18, 2019

Theme: [Environment](#)

In-depth Report: [Climate Change](#)

According to Hans Joachim Schellnhuber, chief climate advisor to the European Union, "We're simply talking about the very life support system of this planet". As fascism and the horror of murderous hate crimes are on the rise, governments are presiding over runaway climate change leading toward mass extinctions of species, costing the lives of billions and the demise of much of nature, while children are protesting against the betrayal of their future.

Evidence based on early climates and on current global warming requires revision of the mostly-linear IPCC climate change trajectories proposed for the 21st to 23rd centuries (Figure 3). The polar ice sheets, acting as thermostats of the climate, are melting at an accelerated rate.

Polar temperatures have been rising at twice the rate of lower latitude zones, weakening the jet stream and the Arctic boundary, which are becoming increasingly undulated (Figure 4). This allows cold air masses to breach the boundary as they move southward, as happened recently in North America and Europe, while warm air masses migrate northward. As the large ice sheets are melting large pools of cold ice melt water are forming in the North Atlantic Ocean south and east of Greenland (Rahmstorf et al. 2015) (See [this](#)). The AMOC (Atlantic Meridional Ocean Circulation) is slowing down and the probability of future transient freeze event/s ([stadials](#)) lasting few decades or longer ([Hansen et al. 2016](#)) is increasing. The juxtaposition of polar-derived freezing fronts and tropics-derived warm air masses leads to a rise in extreme weather events (Figure 1).




Figure 1. Tipping points in the Earth system (Lenton et al., 2008) (see [this](#)) Creative Commons BY-ND 3.0 DE license.

More than 30 years since Professor James Hansen, then NASA's chief climate scientist, presented a stern warning to the US Senate regarding the existential risk posed by global warming (see [this](#)), the consequences of the ignorance, criminal denial and pro-carbon ideology by vested interests and their accomplices among the political classes and in the media, are upon us, as the climate system is shifting into a dangerous uncharted territory. While the Earth as a whole continues to heat, transient temperature polarities between warming land masses and Arctic-derived cold air masses lead to extreme weather events (see [this](#)).

Arctic air temperatures for 2014-2018 have exceeded all previous records since 1900. According to NOAA, Arctic warming has led to a loss of 95 percent of its oldest sea ice over the past three decades. Reports of the International Panel of Climate Change (IPCC), based on thousands of peer reviewed science papers and reports, offer a confident documentation of past and present processes in the atmosphere. On the other hand when it comes to estimates of future ice melt and sea level rise rates, the IPCC models are subject to a number of uncertainties. This includes the difficulty in quantifying amplifying feedbacks from land and water, ice melt rates, linear versus irregular temperature trajectories, sea level rise rates, methane release rates, the role of fires and the observed onset of transient freeze events.




Figure 2. Atmospheric carbon rise rates and global warming events: a comparison between current global warming, the Paleocene-Eocene Thermal Event (PETM) and the last Glacial Termination.

Linear to curved temperature trends portrayed by the IPCC to the year 2300 are rare in the paleo-climate record, where abrupt warming and cooling are common during both glacial and interglacial periods. At +4 degrees Celsius above pre-industrial temperature, projected by the IPCC for the end of the 21st century, life on Earth could be depleted to levels such as existed in the wake of previous mass extinctions of species (see [this](#)).

There would be no smooth transition toward +4 degrees Celsius but irregular heating including a series of extreme weather events and transient temperature reversals induced by the flow of cold ice melt water from the melting glacial sheets into the oceans. Hansen et al. (2016) (see [this](#)) used paleoclimate data and modern observations to estimate the effects of ice melt water from Greenland and Antarctica, with cold low-density meltwater capping warmer subsurface ocean water. Ice mass loss would raise sea level by several and later tens of meters in an exponential rather than a linear response. Sharp drops in temperature, reflecting freeze events in the Atlantic Ocean and the sub-Antarctic Ocean and their surrounds, would reach -2°C for several decades (Figure 3).




Figure 3. Global surface-air temperature to the year 2300 in the North Atlantic and Southern Oceans, including stadial freeze events as a function of Greenland and Antarctic ice melt doubling time (Hansen et al. 2016) (see [this](#))

These projections differ markedly from IPCC models which portray long term ice melting values raising sea levels to less than 1.0 meters by the end of the 21st century (see [this](#)), an estimate difficult to reconcile with satellite gravity-based mass loss estimates by Rignot et al. (2011) (see [this](#)) and others.

With the breach of the Arctic boundary (Figure 4) the world's climate is moving into uncharted territory, with significant implications for the planning of future adaptation efforts, including preparations for sea level rise and for deep freeze events in parts of Western Europe and eastern North America. As the Earth warms the increase in temperature contrasts across the globe, and thereby an increase in storminess and extreme weather events, as occur at present, need to be considered when planning adaptation measures. These would include preparation of coastal defenses from sea level rise and construction of channels and pipelines from flooded regions to drought-stricken zones. In Australia this should include construction of water pipelines and channels from the flooded north to the Murray-Darling basin.

Since many in authority do not accept, or only pay lip service to, climate science, it is a good question whether governments would be investing in adaptation measures in time. In particular no plans appear at hand for draw-down of CO₂– the one measure which could potentially arrest global warming. In this regard the reluctance to date to undertake meaningful mitigation measures does not bode well.

The powers to be are now presiding over the greatest calamity that has ever befallen on humanity and on much of nature.




Figure 4. The undulating jet stream allows penetration of cold air masses from the Arctic southward and warm air masses into the Arctic northward. (see [this](#))

*

Note to readers: please click the share buttons below. Forward this article to your email lists. Crosspost on your blog site, internet forums. etc.

Dr Andrew Glikson, Earth and Paleo-climate science, Australia National University (ANU) School of Anthropology and Archaeology, ANU Planetary Science Institute, ANU Climate Change Institute, Honorary Associate Professor, Geothermal Energy Centre of Excellence, University of Queensland. He is a frequent contributor to Global Research.

The original source of this article is Global Research
Copyright © [Dr. Andrew Glikson](#), Global Research, 2019

[Comment on Global Research Articles on our Facebook page](#)

[Become a Member of Global Research](#)

Articles by: [Dr. Andrew Glikson](#)

Disclaimer: The contents of this article are of sole responsibility of the author(s). The Centre for Research on Globalization will not be responsible for any inaccurate or incorrect statement in this article. The Centre of Research on Globalization grants permission to cross-post Global Research articles on community internet sites as long the source and copyright are acknowledged together with a hyperlink to the original Global Research article. For publication of Global Research articles in print or other forms including commercial internet sites, contact: publications@globalresearch.ca

www.globalresearch.ca contains copyrighted material the use of which has not always been specifically authorized by the copyright owner. We are making such material available to our readers under the provisions of "fair use" in an effort to advance

a better understanding of political, economic and social issues. The material on this site is distributed without profit to those who have expressed a prior interest in receiving it for research and educational purposes. If you wish to use copyrighted material for purposes other than "fair use" you must request permission from the copyright owner.

For media inquiries: publications@globalresearch.ca