

How Climate Change Could Intensify Plague Among Black-tailed Prairie Dogs in the American West

By [Justice Burnaugh](#)

Global Research, February 20, 2018

[The Ecologist](#) 19 February 2018

Region: [USA](#)

Theme: [Environment](#), [Intelligence](#)

In-depth Report: [Climate Change](#)

Featured image: A black-tail prairie dogs during the drought in New Mexico. (Source: David Eads)

Mention the plague and people naturally think of the Black Death during the 14th Century. Antibiotics have since made the plague a curable disease. But the plague is today sterilising ecosystems in the Western United States. And a warming climate could intensify these outbreaks and expand the plague's range farther north and into higher altitudes. JUSTICE BURNAUGH investigates

The sun's heat bears down on a desolate prairie dog colony in America's Great Plains. Small mounds of dirt mark the entrances to an intricate network of tunnels below the surface.

The weeds that the prairie dogs grazed on have reclaimed the soil. A sign reads, "Caution: Prairie Dogs Have Plague. Keep People and Pets in Vehicles."

This is the grim discovery for many biologists returning to study colonies that were thriving only weeks ago. These are the Dark Ages for prairie dogs in the American West.

A new vaccine

Considered a keystone species, black-tailed prairie dogs are an essential part of the American Grasslands. Their tunnels help fertilize the soil and provide shelter for nesting birds.

Many predators rely on prairie dogs as a food source. But the plague's 100 percent mortality rate is shattering this delicate balance.

The plague evolved in Central Asia and made landfall in the United States in 1900 when trade ships from Hong Kong carrying infected rats docked in San Francisco.

The plague spread rapidly throughout the American West where native species like prairie dogs had no immunity to the exotic disease.

Disease researcher Dan Tripp is testing a new vaccine that protects prairie dogs against the plague. The vaccine is fed to the prairie dogs using pellets that are scattered across the colonies. The problem is getting to the prairie dogs before the plague does.

Fleas and rodents

“If you wait until after the plague hits to start vaccinating prairie dogs, then you’re too late,” Tripp said. “By the time you start noticing fewer animals, that’s just the tip of the iceberg of what’s going on below the surface.”

Climate change influences the frequency and severity of plague cycles. Rising temperatures throughout the American West have led to severe droughts and wildfires in some regions and devastating floods in others. This sporadic climate could impede vaccination efforts.

Tripp explained that as the climate wavers between extremes of above average rainfall and droughts, it becomes more difficult for scientists to predict and prevent plague outbreaks.

To understand how climate and plague outbreaks are linked, some scientists are finding their answers with the fleas — the culprits for spreading the plague from one host to another.

If there are more fleas, the chances that one of them is carrying the plague are greater. A popular theory argues that plague outbreaks follow wetter and cooler years, when it’s assumed that fleas and rodents are more abundant.

Wetter times

But new research shows that fleas increase during droughts, meaning that a warmer, drier climate could increase the severity of plague outbreaks in the American West.

Wildlife ecologists David Eads and Dean Biggins counted the amounts of fleas on prairie dogs during wet and dry years to understand how climate affects flea abundance.

The common assumption is that plague outbreaks are more likely during wet years when there are more fleas. To their surprise, they discovered that during a drought in New Mexico the amount of fleas on prairie dogs increased by nearly 200 percent.

“When you have a drought period, flea numbers might escalate on prairie dogs,” Biggins told *The Ecologist*. “And as fleas get more abundant, this could increase the circulation of plague.”

“If we have more drought cycles intermixed with wetter times, which could happen with climate change, you could have more plague circulation.”

Abuzz with speculations

In their [study](#), Eads and Biggins conclude that a lack of food and water during droughts might weaken prairie dogs’ defenses against parasites that spread the plague.

They observed that the fatter, healthier prairie dogs had fewer fleas compared to others that were starving. The study explained that female fleas that feed on starving rodents tend to produce more eggs and larvae.

In light of climate models that forecast frequent droughts, prairie dog colonies in the American West could be at greater risk of plague outbreaks.

When the plague arrived in the American West in 1900, newspapers were abuzz with

speculations about where the plague would hit next.

People knew that fleas spread the plague, but those living in the frigid highlands of the Rocky Mountains believed that the fleas could never survive at such high altitudes. Mountain towns in the American West seemed safely out of reach.

Plague transmission

As the climate continues to warm, northern and high altitude regions that were too cold will become tolerable for fleas and other parasites.

A similar trend is seen with the [northward spread of ticks in America](#), leading to more outbreaks of Lyme disease in the central United States and Canada.

On the Hawaiian island of Kaua'i, [climate change is accelerating the spread of mosquito-borne diseases](#) like malaria into ecosystems in higher elevations, where cooler temperatures are no longer a defense against infectious diseases.

Megan Friggens, an ecologist for the United States Forest Service, said that plague could spread into northern regions of the American West where warming temperatures allow fleas to survive and breed longer.

This increases the amount of fleas and the chances for plague transmission. But Friggens explained that as northern climates become more tolerable for the plague, southern states might become too hot for the plague to survive.

“This heat limitation may actually cause this plague epicenter to shift to the north,” she said. “It may get too hot for fleas in the south, so that would shorten the window for plague transmission in southern regions. You may actually see more plague in northern states because fleas will have a longer [breeding] season.”

*

Justice Burnaugh is a freelance journalist living in the Colorado Rockies. He's been interested in the plague since an outbreak struck a prairie dog colony near his home.

The original source of this article is [The Ecologist](#)
Copyright © [Justice Burnaugh](#), [The Ecologist](#), 2018

[Comment on Global Research Articles on our Facebook page](#)

[Become a Member of Global Research](#)

Articles by: [Justice Burnaugh](#)

not be responsible for any inaccurate or incorrect statement in this article. The Centre of Research on Globalization grants permission to cross-post Global Research articles on community internet sites as long the source and copyright are acknowledged together with a hyperlink to the original Global Research article. For publication of Global Research articles in print or other forms including commercial internet sites, contact: publications@globalresearch.ca

www.globalresearch.ca contains copyrighted material the use of which has not always been specifically authorized by the copyright owner. We are making such material available to our readers under the provisions of "fair use" in an effort to advance a better understanding of political, economic and social issues. The material on this site is distributed without profit to those who have expressed a prior interest in receiving it for research and educational purposes. If you wish to use copyrighted material for purposes other than "fair use" you must request permission from the copyright owner.

For media inquiries: publications@globalresearch.ca