

Melting Ice Sheets and Weakened Polar Fronts: Onset of Climate Tipping Points

By <u>Dr. Andrew Glikson</u> Global Research, February 18, 2019 Theme: <u>Environment</u> In-depth Report: <u>Climate Change</u>

The state of the climate constitutes a confluence of multiple processes, the primary factors being solar insolation and the greenhouse gas composition of the atmosphere. Rising temperatures trigger multiple secondary feedback processes, including decrease in the CO2 solubility in warming parts of the oceans, melting of the large parts of the ice sheets cooling of parts of the oceans by in-flow of ice melt water, release of methane from melting permafrost and methane clathrates and hydrates, desiccated vegetation, bush fires and other feedbacks.

As the Earth continues to heat, transient temperature reversals (stadials) accentuate temperature polarities between warming land and ocean regions cooled by the inflow of ice melt from the ice sheets, as observed south and east of Greenland and off Western Antarctica (Hansen et al. 2016) (Figure 1). Increased polar temperatures, rising twice as fast as intermediate and tropical zones, weaken and undulate the jet stream which defines the polar boundary (Figure 2), allowing cold air masses to move out and warm air masses to move in, further heating the Arctic.

The temperature contrast between migrating warm and cold air masses enhances the intensity of extreme weather events. Analogies are made with Pleistocene and early

Holocene, (2.6-0.01.10⁶ years ago) where peak interglacial temperatures were succeeded by

transient freeze events (stadials), such as the *Younger Dryas* and the 8.5.10³ years-old Laurentide ice melt, attributed to cold ice melt flow into the North Atlantic Ocean and North American lakes (Lake Agasiz). The evidence raises questions regarding the mostly linear to

curved IPCC model trajectories proposed for the 21st century and beyond. Already large pools of cold ice melt are formed in the ocean south and east of Greenland (Rahmstorf et al. 2015) and north of west Antarctica (Figure 1) and the AMOC (Atlantic Meridional Ocean Circulation) and the jet stream are weakening. Comparisons of current warming with past

climates of $+1^{\circ}$ C relative to the early 19^{th} century indicate extreme weather events, including in the early Holocene Optimum period (~10,000-8500 years ago) and the Eemian interglacial (130,000 – 115,000 years ago) (Roverea et al. 2017). The probability of a future transient freeze event (stadial) triggered by the flow of cold icemelt water into the North Atlantic and sub-Antarctic oceans bears major implications for modern and future climate change trends, including a rise in extreme weather events due to a growing contrast between cooling oceans and warming continents and between polar-derived and tropics-derived air masses of contrasted temperatures. This needs to be taken into account in planning adaptation efforts.



Figure 1.Modelled 2055-2100 surface-air temperature to +1.19°C above 1880-1920 (AIB modelmodified forcing, ice melt to 1 meter sea lelvel rise) (Hansen et al. 2016)¹



Figure 2. The weakening and increasing undulation of the polar boundary (see this)

Whereas reports of the International Panel of Climate Change (IPCC²), based on thousands of peer reviewed science papers and reports, offer a confident documentation of past and present processes in the atmosphere³, including future model projections (Figure 3), when it comes to estimates of future ice melt and sea level change rates, however, these models contain a number of significant departures from observations based on the paleoclimate record. This includes climate change feedbacks from land and water, ice melt rates, temperature trajectories, sea level rise rates, methane release rates, the role of fires, and observed onset of transient stadial (freeze) events⁴. Early stages of stadial event/s are

manifest by the build-up of a large pools of cold ice melt water in the North Atlantic Ocean south of Greenland and along the fringes of the Antarctic continent (Figure 1).



Figure 3. IPCC model of the time series of the global annual mean surface air temperatureanomalies relative to 1986–2005 from CMIP5 (Coupled Model Inter- comparison Project) Projections are shown for

each multi model mean (solid lines)⁵

Hansen et al. (2016) (Figure 1) used paleoclimate data and modern observations to estimate theeffects of ice melt water from Greenland and Antarctica, showing cold lowdensity meltwater tend to cap increasingly warm subsurface ocean water, affecting an increase in ice shelf melting, accelerating ice sheet mass loss (Figure 4) and slowing deep water formation (Figure 5). Ice mass loss would raise sea level by several meters as an exponential rather than linear response, with doubling time of ice loss of 10, 20 or 40 years yielding multi-meter sea level rise in about 50, 100 or 200 years.

Linear to curved temperature trends portrayed by the IPCC to the year 2300 (Figure 3) are rare in the Pleistocene paleo-climate record, where abrupt warming and cooling dominate

during glacial periods (Dansgaard-Oeschger cycles; Ganopolski and Rahmstorf 2001⁶;

Camille and Born, 2019⁷) and interglacial (Cortese et al. 2007⁸) periods.Hansen et al.'s (2016) model includessharp drops in temperature, reflecting stadial freezing events in the

Atlantic Ocean and the sub-Antarctic Ocean and their surrounds, reaching -2°C over several decades (Figure 6).



Figure 4.Greenland and Antarctic ice mass change. GRACE data are extension of Velicogna etal. (2014)⁹gravity data. MBM (mass budget method) data are from Rignot et al. (2011)¹⁰. Redcurves are gravity data for Greenland and Antarctica; small Arctic ice caps and ice shelf melt add to freshwater

input (Hansen et al. 2016)¹¹



Figure 5. The AMOC (Atlantic Mid-Ocean circulation) at 28°N in simulations (i.e., includingfreshwater injection of 720 Gt year-1 in 2011 around Antarcti ca, increasing with a 10-year doubling time, and half

that amount around Greenland). (b) SST (°C) in the North Atlantic region (44–60 $\,^\circ$ N, 10–50 $\,^\circ$ W).

Temperature and sea level rise relations during the Eemian interglacial¹² about 115-130 kyr ago, when temperatures were about $+1^{\circ}$ C or higher than during the late stage of the Holocene, and sea levels were +6 to +9 m higher than at present, offer a possible analogy for present developments. During the Eemian overall cooling of the North Atlantic Ocean and parts of the West Antarctic fringe ocean due to ice melt led to increased temperature polarities and to storminess (Roverea etal. 2017; Kaspar et al. 2007)¹³, underpinning the danger of global temperature rise to $+1.5^{\circ}$ C. Accelerating ice melt and nonlinear sea level



Figure 6. Global surface-air temperature to the year 2300 in the North Atlantic and SouthernOceans, including stadial freeze events as a function of Greenland and Antarctic ice melt doubling time

The development of large cold water pools south and east of Greenland (Rahmstorf et al.

2015¹⁴) and at the fringe of West Antarctica (Figure 1) signify early stages in the development of a stadial freeze, consistent with the decline in the Atlantic Meridional Ocean Circulation (AMOC) (Figure 5). These projections differ markedly from model trends of the IPCC (Figure 3), which portray long term ice melt (Ahmed N 2018)¹⁵. The IPCC

(2016)¹⁶ states: A key question is whether ice-dynamical mechanisms could operate which would enhance ice discharge sufficiently to have an appreciable additional effect on sea level rise". This statement is difficult to reconcile with studies by Rignot et al. (2011), reporting that in 2006the Greenland and Antarctic ice sheets experienced a combined mass

loss of 475 \pm 158 Gt/yr, equivalent to 1.3 \pm 0.4 mm/yr sea level rise."¹⁷ For the Antarctic ice

sheet the IEMB team $(2017)^{18}$ states the sheet lost 2,720 ± 1,390 billion tonnes of ice between 1992 and 2017, which corresponds to an increase in mean sea level of 7.6 ± 3.9 millimeter.

A non-linear climate warming trend, including stadial freeze events, bears significant implications for planning future adaptation efforts, including preparations for transient deep freeze events in parts of Western Europe and eastern North America, for periods lasting several decades (Figure 6), as well as coastal defenses against enhanced storminess arising from increased temperature contrasts between the cooled regions and warm tropical latitudes.

According to NOAA¹⁹ Arctic surface air temperatures continue to warm at twice the rate relative to the rest of the globe, leading to a loss of 95 percent of its oldest ice over the past three decades. Arctic air temperatures for 2014-18 have exceeded all previous records since 1900 and are driving broad changes within the Arctic as well the sub-Arctic through weakening of the jet stream which separates the Arctic from warmer climate zones. The recent freezing storms in North America represent penetration of cold air masses through a weakening and increasingly undulating jet stream barrier (Figures 2 and 7). This weakening also allows warm air masses to move northward, further warming the Arctic and driving further ice melting. The freezing storms in North America are cheering climate denialists who refuse to discriminate between the climate and the weather. As the Earth continues to heat and cold air masses breach the Arctic boundary and move southward, temperature

contrasts between polar and subpolar climate zones decrease, further weakening the polar divide. At the same time temperature contrasts between Arctic-derived cold air masses and subtropical zones result in an increase in the intensity and frequency of extreme weather events.



Figure 7. The weakened undulating Jet stream bounding the polar vortex. Red represents the fastest air flow (Berwyn 2016) 20



Figure 8. The North American and Siberian freeze event 30 January 2019 (NOAA GlobalForecast system

The role of feedbacks from land and water, estimates of future ice melt rates, sea level rise rates, methane release rates, the role of fires in enhancing atmospheric CO_2 , and the already observed onset of temporary freeze events need to be quantified. As the Earth warms, the increase in temperature contrasts across the globe, and thereby an increase in storminess and extreme weather events, occurring at present, need to be taken into account when planning adaptation measures, including preparation of coastal defenses, construction of channel and pipelines from heavy precipitation zones to draught zones. In Australia this should include construction of water pipelines and channels from the flooded north to parched regions such as the Murray-Darling basin.

Imminent climate risks

Climate model projections for the 21st to 23rd centuries need to take paleoclimate evidence more fully into account, including the transient stadial effects of ice melt water flow into the oceans and amplifying feedbacks of global warming from land and oceans. The paleoclimate record indicates that over the last 800,000 years peak interglacial temperatures were consistently succeeded by temporary freeze events, attributed to the flow of cold ice melt water flow into the North Atlantic Ocean. Radiative forcing²², increasing with concentration of atmospheric greenhouse gases and rising by about 0.04 Watt/m²/year over the last 50 years²³, totaled by more than 2 Watt/m², equivalent to ~3.0°C (~1.5°C per W/m ²)²⁴. The rise of mean global temperatures to date by 0.9°C since 1880 ²⁵ therefore represents lag effect, pointing to potential temperature rise by approximately two degrees Celsius. Climate change trajectories would be highly irregular as a result of stadial events affected by flow of ice melt water into the oceans. Whereas similar temperature fluctuations

and stadial events occurred during past interglacial periods (Cortese etal. 2007²⁶; Figure 9),

when peak temperature fluctuations were close to $+/1^{\circ}$ C, further rises intemperature in future would enhance the intensity and frequency of extreme weather events, entering uncharted territory, rendering large parts of the continents uninhabitable.



Figure 9.A.Evolution of sea surface temperatures in 5 glacial-interglacial transitions recordedin

ODP-1089 at the sub-Antarctic Atlantic Ocean. Lower grey lines – δ^{18} O measured on Cibicidoides plankton; Black lines – sea surface temperature. Marine isotope stage numbers are indicated on top of diagrams. Note the stadial temperature drop events following interglacial peak temperatures,

analogous to the *Younger Dryas*preceding the onset of the Holocene (Corteseet al. 2007²⁷).B. Mean temperatures for the late Pleistocene and early Holocene.

*

Note to readers: please click the share buttons below. Forward this article to your email lists. Crosspost on your blog site, internet forums. etc.

Dr Andrew Glikson, Earth and Paleo-climate science, Australia National University (ANU) School of Anthropology and Archaeology, ANU Planetary Science Institute, ANU Climate Change Institute, Honorary Associate Professor, Geothermal Energy Centre of Excellence, University of Queensland. He is a frequent contributor to Global Research.

Sources

Arctic Report Card (2018) Tracking recent environmental changes relative to historical records https://www.arctic.noaa.gov/Report-Card

Berwyn B (2016) Wobbly Jet Stream Is Sending the Melting Arctic into 'Uncharted territory. Inside Climate News

https://insideclimatenews.org/news/08062016/greenland-arctic-record-melt-jet-stream-wobb ly-global-warming-climate-change

Camille Li, Born A (2019) Coupled atmosphere-ice-ocean dynamics in Dansgaard-Oeschger events Quaternary Science Reviews 203, 1-20.

https://www.sciencedirect.com/science/article/pii/S0277379118305705

Climate Council (1918) The good the b ad and the ugly: limiting temperature rise to 1.5°C. https://www.climatecouncil.org.au/resources/limiting-temperature-rise/

Cortese G, Abelmann A, Gersonde A (2007) The last five glacial-interglacial transitions: A high-resolution 450,000-year record from the sub-Antarctic Atlantic. Paleogeography and Paleoclimatology (22) Part 4.

https://www.researchgate.net/publication/228524417_The_last_five_glacial-interglacial_trans itions_A_high-resolution_450000-year_record_from_the_subantarctic_Atlantic

Easterbrook S (2019) New IPCC Report (Part 6). Azimuth. https://johncarlosbaez.wordpress.com/2014/04/16/what-does-the-new-ipcc-report-say-aboutclimate-change-part-6/

Francis J (2019) How frigid polar vortex blasts are connected to global warming: The National Weather Service is warning of brutal, life-threatening conditions. Salon, January 2019.

https://www.salon.com/2019/01/31/how-frigid-polar-vortex-blasts-are-connected-to-global-w arming_partner/

Ganopolski A, Rahmstorf S. (2001) Rapid changes of glacial climate simulated in a coupled climate model. Nature 409 (6817)153-8. https://www.ncbi.nlm.nih.gov/pubmed/11196631

Hansen J. et al. (2016) Ice melt, sea level rise and superstorms: evidence from paleoclimate data, climate modeling, and modern observations that 2°C globa I warming could be dangerous. Atmos. Chem. Phys. 16, 3761-3812.

New IPCC Report (Part 6)

IMBIE Team (2017) Mass balance of the Antarctic Ice Sheet from 1992 to 2017. The IMBIE team.

https://www.nature.com/articles/s41586-018-0179-y.epdf?referrer_access_token=S5Y_R7foK De_0LTC1ePHNRgN0jAjWel9jnR3ZoTv0PBEKqWHTwARrIrR4OxoHFd5WZGhA0FX8FPbkdWIZL YWSZXdrY6PsBEIhQw8kfzqY8CzRUyWaogOmRIMtURwKL_LY17cUVdIgmtWLaRk_EWhFIL0JdJy awITzJhU3y8fPcoosWQQMgEN2fv3kQx_S8JT4BLn4bheLaGZaYfD6J64pzwLO1V5h5TxsI6J4qUi mPnWHm2Ax0DoQjYvfEgChVqY1nI8d3M_kRuObyJedPw%3D%3D&tracking_referrer=www.ab c.net.au

IPCC (2018) Global warming of 1.5oC. https://www.ipcc.ch/sr15/

IPCC Working Group I (2018) The Scientific Basis. https://archive.ipcc.ch/ipccreports/tar/wg1/416.htm; https://www.ipcc.ch/working-group/wg1/?idp=408

Kaspar, F, Spangehl T, and Cubasch U (2007). Northern hemisphere winter storm tracks of the Eemian interglacial and the last glacial inception. Clim. Past 3, 181–192

Nafeez Ahmed (2018) |The UN's Devastating Climate Change Report Was Too Optimistic. Mother board. https://motherboard.vice.com/en_us

Rahmstorf S. et al. (2015) Exceptional twentieth-century slowdown in Atlantic Ocean overturning circulation. Nature Climate Change volume 5:475-480

Rignot E et al. (2011) Acceleration of the contribution of the Greenland and Antarctic ice sheets to sea level rise Acceleration of the contribution of the Greenland and Antarctic ice sheets to sea level rise. Geophysical Res Lett 38 (5).

Roverea A et al. (2017) Giant boulders and Last Interglacial storm intensity in the North Atlantic. Proc. Am Acad Sci 114 (46) 12144-12149.

Velicogna I, Sutterley T. C. vanden Broeke M. R. (2014) Regional acceleration in ice mass loss from Greenland and Antarctica using GRACE time-variable gravity data. Geophysical Res Lett 41 (22) 8130-8137 https://agupubs.onlinelibrary.wiley.com/doi/10.1002/2014GL061052

Notes

1. https://www.atmos-chem-phys.net/16/3761/2016/

2. IPCC (2018) Global warming of 1.5 degrees. <u>https://www.ipcc.ch/sr15/</u>

3.

https://www.climatecouncil.org.au/resources/limiting-temperature-rise/?gclid=EAIaIQobChMlt-Wq7-KM4 AIVVQ4rCh0WSQ69EAAYASAAEgLhp_D_BwE

4. https://www.atmos-chem-phys.net/16/3761/2016/acp-16-3761-2016.pdf

5.

https://johncarlosbaez.wordpress.com/2014/04/16/what-does-the-new-ipcc-report-say-about-climate-cha nge-part-6/

6. https://www.ncbi.nlm.nih.gov/pubmed/11196631

7. https://www.sciencedirect.com/science/article/pii/S0277379118305705

8. https://agupubs.onlinelibrary.wiley.com/doi/full/10.1029/2007PA001457

9. https://agupubs.onlinelibrary.wiley.com/doi/10.1002/2014GL061052

10. https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2011GL046583

11. https://www.atmos-chem-phys.net/16/3761/2016/acp-16-3761-2016.pdf

12. https://www.britannica.com/science/Eemian-Interglacial-Stage

13.

http://moraymo.us/wp-content/uploads/2018/03/Rovereetal_PNAS_2017.pdf;https://www.clim-past.net/3/ 181/2007/cp-3-181-2007.pdf

14. https://www.nature.com/articles/nclimate2554

15.

https://motherboard.vice.com/en_us/article/43e8yp/the-uns-devastating-climate-change-report-was-too-optimistic

16. https://archive.ipcc.ch/ipccreports/tar/wg1/416.htm

17. https://agupubs.onlinelibrary.wiley.com/doi/full/10.1029/2011GL046583

18. https://www.nature.com/articles/s41586-018-0179-y

19. https://www.arctic.noaa.gov/Report-Card

20. https://insideclimatenews.org/news/08062016/greenland-arctic-record-melt-jet-stream-wobbly-globa l-warming-climate-change

21. https://www.salon.com/2019/01/31/how-frigid-polar-vortex-blasts-are-connected-to-global-warming_partner/

22. Radiative forcing - the difference between incoming radiation and radiation reflected back to space

- 23. http://www.columbia.edu/~jeh1/mailings/2018/20181206_Nutshell.pdf
- 24. https://arxiv.org/abs/0804.1126

25. https://climate.nasa.gov/vital-signs/global-temperature/

26. https://agupubs.onlinelibrary.wiley.com/doi/full/10.1029/2007PA001457

27. https://agupubs.onlinelibrary.wiley.com/doi/full/10.1029/2007PA001457

The original source of this article is Global Research Copyright © <u>Dr. Andrew Glikson</u>, Global Research, 2019

Comment on Global Research Articles on our Facebook page

Become a Member of Global Research

Articles by: Dr. Andrew Glikson

Disclaimer: The contents of this article are of sole responsibility of the author(s). The Centre for Research on Globalization will not be responsible for any inaccurate or incorrect statement in this article. The Centre of Research on Globalization grants permission to cross-post Global Research articles on community internet sites as long the source and copyright are acknowledged together with a hyperlink to the original Global Research article. For publication of Global Research articles in print or other forms including commercial internet sites, contact: publications@globalresearch.ca

www.globalresearch.ca contains copyrighted material the use of which has not always been specifically authorized by the copyright owner. We are making such material available to our readers under the provisions of "fair use" in an effort to advance a better understanding of political, economic and social issues. The material on this site is distributed without profit to those who have expressed a prior interest in receiving it for research and educational purposes. If you wish to use copyrighted material for purposes other than "fair use" you must request permission from the copyright owner.

For media inquiries: publications@globalresearch.ca