

# The Fading Home of Homo 'Sapiens'. Extreme Weather Events

By [Dr. Andrew Glikson](#)

Global Research, August 15, 2019

Theme: [Environment](#)

In-depth Report: [Climate Change](#)

*In front of our eyes green forests are blackened by fire, draughts turn grassy planes to brown semi-deserts, fertile fields are flooded by muddy water, white snow and ice melt to pale blue water and clear blue skies turn grey with aerosols.*


\*\*\*

The inhabitants of planet Earth are in the process of destroying the habitability of their world through the perpetration of the largest mass extinction of species since [66 million years](#) ago when a large asteroid impacted Earth and 55 million years since the Paleocene-Eocene Thermal Maximum ([PETM](#)) reaching +5 to +8°C. The late Holocene-Anthropocene climate change represents an unprecedented event, triggering a fast shift in climate zones and a series of extreme weather events, with consequences for much of nature and civilization. The changes are manifest where green forests are blackened by fire, draughts are turning grassy planes to brown semi-deserts, brilliant white snow and ice caps are melting into pale blue water and clear blue skies turn grey due to aerosols and jet contrails, most particularly in the northern hemisphere. Unless effective efforts are undertaken at CO<sub>2</sub>down-draw, the consequence would include demise of much of nature and a collapse of human civilization.

## 1. The scorched Earth

The transfer of hundreds of billions of tonnes of carbon from the Earth crust, the residues of ancient biospheres, to the atmosphere and oceans, condemning the bulk of life through the most extreme shift in the composition of the atmosphere and ocean Earth has experienced since 55 million years ago, with changes taking place in front of our eyes. Since the industrial revolution, about [375 billion tonnes of carbon](#) (or 1,374 billion tonnes CO<sub>2</sub>) have been emitted by humans into the atmosphere. The consequences are everywhere, from mega-droughts, to heat waves, fires, storms and floods. With atmospheric CO<sub>2</sub>-equivalent rising [above 500 ppm](#) and mean temperatures by more than 1.5°C (Figure 1) look no further than the shift in climate zones, displayed for example on [maps of the expanding wet tropical zones](#), drying sub-tropical latitudes and polar-ward migration of temperate climate zone. The ice sheets and sea ice are melting, huge fires overtake Siberia, the [Sahara is shifting northward](#), large parts of southern Europe are suffering from draughts, heat waves and fires, the Kalahari Desert dunes are [shifting](#) and much of [southern Australia is affected by warming and draughts](#). This is hardly compensated by a minor increase in precipitation and greening such as at the southern fringes of the Sahara Desert and parts of northern Australia.

Induced by anthropogenic carbon emissions reaching 37.1 billion tonnes CO<sub>2</sub> in 2018 and their amplifying feedbacks from land and oceans, ranging from where the US produces 16.5 tonnes CO<sub>2</sub> per capita per year, to 35.5 tonnes CO<sub>2</sub> per capita per year from Saudi Arabia's and 44 tons export per capita [per year](#) from Australia. The inexorable link between these emissions and the unfolding disaster is hardly mentioned by mainstream political classes and the media.



[Figure 1](#) (A) Growth of CO<sub>2</sub>-equivalent level and the annual greenhouse gas Index (AGGI<sup>[1]</sup>). Measurements of CO<sub>2</sub> to the 1950s are from (Keeling et al., 1958) and air trapped in ice and snow above glaciers. Pre-1978 changes are based on ongoing measurements of all greenhouse gases. Equivalent CO<sub>2</sub> amounts (in ppm) are derived from the relationship between CO<sub>2</sub> concentrations and radiative forcing from all [long lived gases](#); (B) showing how much warmer each month of the [GISTEMP](#) data is than the annual global mean. For July (2019) temperatures rose by about +1.5°C.

## 2. Migrating climate zones

As the globe warms, to date by a mean of near [~1.5°C](#), or [~2.0°C](#) when the masking effects of sulphur dioxide and other aerosols are considered, and by a mean of [~2.3°C](#) in the polar regions, the expansion of [warm tropical latitudes](#) and the pole-ward [migration of subtropical and temperate climate zones](#) (Figure 2) ensue in large scale draughts such as parts of inland [Australia](#) and [southern Africa](#). A similar trend is taking place in the northern hemisphere where the Sahara desert is expanding northward, with consequent heat waves across the [Mediterranean and Europe](#).

In southern Africa “[Widespread shifts in climate regimes](#)” are projected, of which the southern and eastern expansion of the hot desert and hot steppe zones is most prominent. From occupying 33.1 and 19.4 % of southern Africa under present-day climate, respectively, these regions are projected to occupy between 47.3 and 59.7 % (hot desert zone) and 24.9 and 29.9 % (hot steppe zone) of the region in a future world where the mean global temperature has increased by  $\sim 3^{\circ}\text{C}$ .

Closely linked to the migration of climate zones is the southward drift of Antarctic- sourced cold moist fronts which sustain seasonal rain in south-west and southern Australia. A feedback loop has developed where deforestation and decline in vegetation in southern parts of the continent result in the rise of thermal plumes of dry air masses that deflect the western moist fronts further to the southeast.




Figure 2. Köppen-Geiger global Climate zones classification map

An updated Köppen–Geiger climate map<sup>[1]</sup>



- **2Group A: Tropical/megathermal climates**
  - **2.1Af: Tropical rainforest climate**
  - **2.2Am: Tropical monsoon climate**
  - **2.3Aw/As: Tropical wet and dry or savanna climate**
- **3Group B: Dry (desert and semi-arid) climates**
  - **3.1BW: Arid Climate**
  - **3.2BS: Semi-arid (Steppe) Climate**
- **4Group C: Temperate/mesothermal climates**
  - **4.1Csa: Mediterranean hot summer climates**
  - **4.2Csb: Mediterranean warm/cool summer climates**
  - **4.3Csc: Mediterranean cold summer climates**
  - **4.4Cfa: Humid subtropical climates**
  - **4.5Cfb: Oceanic climate**
  - **4.6Cfb(2): Subtropical Highland climates with uniform rainfall**
  - **4.7Cfc: Subpolar oceanic climate**
  - **4.8Cwa: Dry-Winter Humid Subtropical**
  - **4.9Cwb: Dry-Winter Subtropical Highland**
  - **4.10Cwc: Dry-Winter Subpolar Oceanic climate**
- **5Group D: Continental/microthermal climates**
  - **5.1Dfa/Dwa/Dsa: Hot summer continental climates**
  - **5.2Dfb/Dwb/Dsb: Warm summer continental or hemiboreal climates**
  - **5.3Dfc/Dwc/Dsc: Subarctic or boreal climates**
  - **5.4Dfd/Dwd/Dsd: Subarctic or boreal climates with severe winters**
- **6Group E: Polar climates**
  - **6.1ET: Tundra climate**
  - **6.2EF: Ice cap climate**
- **7Ecological significance**

Since 1979 the planet's tropics have been [expanding pole-ward by 56 km to 111 km per decade](#) in both hemispheres, leading one commentator to call this [Earth's bulging waistline](#). Future climate projections suggest this expansion is likely to continue, driven largely by human activities – most notably emissions of greenhouse gases and black carbon, as well as warming in the lower atmosphere and the oceans.”

An analysis of the origin of [Australian draughts](#) suggests, according to both observations and climate models, that at least part of this decline is associated with changes in large-scale atmospheric circulation, including shrinking polar ice and a [pole-ward movement of polar-originated westerly wind spirals](#), as well as increasing atmospheric surface pressure and droughts over parts of southern Australia (Figure 3). Simulations of future climate with this model suggest amplified winter drying over most parts of southern Australia in the coming decades in response to changes in radiative forcing. The drying is most pronounced over southwest Australia, with total reductions in austral autumn and winter precipitation of approximately [40% by the late twenty-first century](#). Thus rainfall in southwestern Australia has declined sharply from about [1965 onward](#), concomitant with the sharp [rise of global temperatures](#).






Figure 3(A) Bureau of Meteorology (BOM) draught map, showing rainfall levels for the southern wet season from April 1 to July 31 in 2019; (B) NASA satellite image displaying a southward deflection of Antarctic-sourced moist cold fronts from southern Australia, a result of (1) southward migration of climate zones; (2) increasing aridity of southern and southwestern Australia due to deforestation; (3) rising hot plumes from warming arid land..

### 3. Extreme weather events

The consequences of the migration of climate zones are compounded by changes in flow patterns of major [river systems](#) around the world, for example in southern and southeastern Asia, including the Indus, Ganges, Brahmaputra and Mekong river basins, the home and bread basket for more than a billion people. With warming, as snow cover declines in the mountainous source regions of rivers, river flows are enhanced, with ensuing floods, in

particular during the Monsoon. For example, in 2010 approximately one-fifth of Pakistan's total land area was affected by floods (Figure 4A), directly affecting about 20 million people, with a death toll close to [2,000](#). And about 700 people in cyclone Isai in Mozambique (Figure 4B, C). Such changes in climate and geography are enhanced once sea level rise increases from the scale of tens of centimeters, as at present, to [meters](#), as predicted to take place later this and next century.

An increasing frequency and intensity of cyclones constitute an inevitable consequence of rising temperatures over warm low pressure cell tracks in tropical oceans, already affecting large populations in the Caribbean and west Pacific island chains, encroaching into continental coastal zones, [China](#), [southeast USA](#), [southeast Africa](#), [India](#), [northern Australia](#), the [Pacific islands](#). According to [Sobel et al. \(2016\)](#)

“We thus expect tropical cyclone intensities to increase with warming, both on average and at the high end of the scale, so that the strongest future storms will exceed the strength of any in the past”.

Likewise increasing temperatures, heat waves and droughts, compounded by deforestation over continents, constitute an inevitable consequence of heat waves and droughts. A prime example is the Siberian forest fires (Figure 5B), covering an area larger than Denmark and contributing significantly to climate change. Since the beginning of the year a total of [13.1 million hectares](#) has burned. Total losses from natural catastrophes on 2018 stated as [US\\$160 billion](#).




Figure 4(A) Pakistan flooding showing the 2010 Indus River spanning well over 10 kilometers, completely filling the river valley and spilling over onto nearby land. Floodwaters have created a lake almost as wide as the swollen Indus that inundates Jhatpat; (B) Before-and-after satellite imagery of Mozambique showing massive flood described as an “inland ocean” up to 30 miles wide following the landfall of Tropical [Cyclone Idai](#), 2019.






Figure 5(A) Global fire zones, NASA. The Earth data fire map accumulates the locations of fires detected by moderate-resolution imaging radiometer ([MODIS](#)) on board the Terra and Aqua satellites over a 10-day period. Each colored dot indicates a location where MODIS detected at least one fire during the compositing period. Color ranges from red where the fire count is low to yellow where number of fires is large;(B)An ecological catastrophe in Russia: wildfires have created over 4 million square km smoke lid over central northern Asia. Big Siberian cities are covered with toxic haze that had already reached Urals.

#### 4. Shrinking Polar ice sheets

Last but not least, major changes in the Polar Regions are driving climate events in the rest of the globe. According to [NOAA](#) Arctic surface air temperatures continued to warm at [twice the rate](#) of the rest of the globe, leading to major thaw at the fringes of the Arctic (Figure 6A) and a loss of [95 percent of its oldest ice](#) over the past three decades. Arctic air temperatures during 2014-18 since 1900 have [exceeded all previous records](#) and are driving broad changes in the environmental system both within the Arctic as well as through the [weakening](#) of the jet stream which separates the Arctic from warmer climate zones. The recent freezing storms in North America represent penetration of cold air masses through an increasingly undulating jet stream barrier, as well as allowing warm air masses to move northward, further warming the Arctic and driving further ice melting (Figure 6B).



According to [Rignot et al. \(2011\)](#) in 2006 the Greenland and Antarctic ice sheets experienced a combined mass loss of  $475 \pm 158$  billion tons of ice per year. IPCC models of future climate change contain a number of departures from the paleoclimate evidence, including the major role of feedbacks from land and water, estimates of future ice melt, sea level rise rates, methane release rates, the role of fires in enhancing atmospheric CO<sub>2</sub>, and the already observed onset of transient freeze events consequent on the [flow of ice melt water into the oceans](#). Ice mass loss would raise sea level on the scale of meters and eventually tens of meters ([Hansen et al. 2016](#)). The development of large cold water pools south and east of Greenland ([Rahmstorf et al. 2015](#)) and at the fringe of West Antarctica, signify early stages in the development of a North Atlantic freeze, consistent with the decline in the Atlantic Meridional Ocean Circulation ([AMOC](#)). As the Earth warms the increase in temperature contrasts across the globe, in particular between warming continental regions and cooling ocean regions, leads to storminess and extreme weather events, which need to be taken into account when planning adaptation measures, including preparation of coastal defenses, construction of channel and pipelines from heavy precipitation zones to draught zones.




Figure 6(A) [Thawing at the fringes of Siberia and Canada](#). Scientists say 2019 could be another *annus horribilis* for the Arctic with record temperatures already registered in Greenland—a giant melting ice sheet that threatens to submerge the world’s coastal areas one day; (B) Weakening and increasing undulation of the [polar vortex](#), allowing penetration of cold fronts southward and of warm air masses northward.




Figure 7 (A) Surface air temperature ( $^{\circ}\text{C}$ ) change in 2055-2060 relative to 1880-1920 according to AIB model + modified forcings and ice melt to 1 meter sea level rise; (B) Surface-air temperature change in 2096 relative to 1880-1920 according to IPCC model AIB adding Ice melt with 10-year doubling of ice melt leading to +5 meters sea level rise; (C) Surface air temperature ( $^{\circ}\text{C}$ ) relative to 1880-1920 for several scenarios taking added ice melt water into account ([Hansen et al. 2016](#))

## Postscript

None of the evidence and projections summarized above appears to form a priority consideration on the part of those in power—in parliaments, in corporations or among the wealthy elites and vested interests. Having to all intents and purposes given up on the [habitability of large parts of the Earth](#) and on the survival of numerous species and future generations—their actions and inactions constitute the ultimate crime against life on Earth.

\*

Note to readers: please click the share buttons above or below. Forward this article to your email lists. Crosspost on your blog site, internet forums. etc.

*Dr Andrew Glikson, Earth and Paleo-climate science, Australia National University (ANU) School of Anthropology and Archaeology, ANU Planetary Science Institute, ANU Climate Change Institute, Honorary Associate Professor, Geothermal Energy Centre of Excellence, University of Queensland. He is a frequent contributor to Global Research.*

Note

[1] The [AGGI index](#) uses 1990 as a baseline year with a value of 1. The index increased every year since 1979.

The original source of this article is Global Research  
Copyright © [Dr. Andrew Glikson](#), Global Research, 2019

---

[Comment on Global Research Articles on our Facebook page](#)

[Become a Member of Global Research](#)

Articles by: [Dr. Andrew Glikson](#)

**Disclaimer:** The contents of this article are of sole responsibility of the author(s). The Centre for Research on Globalization will not be responsible for any inaccurate or incorrect statement in this article. The Centre of Research on Globalization grants permission to cross-post Global Research articles on community internet sites as long the source and copyright are acknowledged together with a hyperlink to the original Global Research article. For publication of Global Research articles in print or other forms including commercial internet sites, contact: [publications@globalresearch.ca](mailto:publications@globalresearch.ca)

[www.globalresearch.ca](http://www.globalresearch.ca) contains copyrighted material the use of which has not always been specifically authorized by the copyright owner. We are making such material available to our readers under the provisions of "fair use" in an effort to advance a better understanding of political, economic and social issues. The material on this site is distributed without profit to those who have expressed a prior interest in receiving it for research and educational purposes. If you wish to use copyrighted material for purposes other than "fair use" you must request permission from the copyright owner.

For media inquiries: [publications@globalresearch.ca](mailto:publications@globalresearch.ca)